(Agrs 11-15 ye old) PHQ-9: Modified

Clinician:

Date:

Name:

	Instructions: How often have you been be past two weeks? For each symptom put a describes how you have been feeling.	othered by in " X" in th	each of the fol ne box beneath	llowing sympton the answer the	oms during the nat best)
			Not At All	Several Days	(2) More Than Half the Days	Nearly Every Day
1.	Feeling down, depressed, irritable, or hopeles	s?				
2.	Little interest or pleasure in doing things?					
3.	Trouble falling asleep, staying asleep, or sleep much?	oing too				
4.	Poor appetite, weight loss, or overeating?					
5.	Feeling tired, or having little energy?					
6.	Feeling bad about yourself – or feeling that you failure, or that you have let yourself or your far	u are a nily				
	down?				LI	
7.	Trouble concentrating on things like school wo reading, or watching TV?	ork,				
8.	Moving or speaking so slowly that other people have noticed? Or the opposite – being so fidgety or restless twere moving around a lot more than usual?	hat you				
·	Thoughts that you would be better off dead, or hurting yourself in some way?					
In the <u>past year</u> have you felt depressed or sad most days, even if you felt okay sometimes? [] Yes [] No						
If you are experiencing any of the problems on this form, how difficult have these problems made it for you to do your work, take care of things at home or get along with other people? [] Not difficult at all [] Somewhat difficult [] Very difficult [] Extremely difficult						
Has there been a time in the <u>past month</u> when you have had serious thoughts about ending your life? [] Yes [] No						
Have you <u>EVER</u> , in your WHOLE LIFE, tried to kill yourself or made a suicide attempt? [] Yes [] No						
2 8 8000	**If you have had thoughts that you would be better off dead or of hurting yourself in some way,					

Office use only: Severity score:

please discuss this with your Health Care Clinician, go to a hospital emergency room or call 911.

Scoring the PHQ-9 modified for Teens

Scoring the PHQ-9 modified for teens is easy but involves thinking about several different aspects of depression.

To use the PHQ-9 as a diagnostic aid for Major Depressive Disorder:

- Questions 1 and/or 2 need to be endorsed as a "2" or "3"
- Need five or more positive symptoms (positive is defined by a "2" or "3" in questions 1-8 and by a "1", "2", or "3" in question 9).
- The functional impairment question (How difficult....) needs to be rated at least as "somewhat difficult."

To use the PHQ-9 to screen for all types of depression or other mental illness:

- All positive answers (positive is defined by a "2" or "3" in questions 1-8 and by a "1", "2", or "3" in question 9) should be followed up by interview.
- A total PHQ-9 score ≥ 10 (see below for instructions on how to obtain a total score) has a good sensitivity and specificity for MDD.

To use the PHQ-9 to aid in the diagnosis of dysthymia:

 The dysthymia question (In the past year...) should be endorsed as "yes."

To use the PHQ-9 to screen for suicide risk:

 All positive answers to question 9 as well as the two additional suicide items MUST be followed up by a clinical interview.

To use the PHQ-9 to obtain a total score and assess depressive severity:

- Add up the numbers endorsed for questions 1-9 and obtain a total score.
- See Table below:

Total Score	Depression Severity
0-4	No or Minimal depression
5-9	Mild depression
10-14	Moderate depression
15-19	Moderately severe depression
20-27	Severe depression